

EVALUATION OF COLD RESISTANCE AND EARLY RIPENING TRAITS IN STERILE FORMS OF TOMATOES

S. SHOKH, L. SHUBENKO, ZH. VDOVYCHENKO

*Белоцерковский национальный аграрный университет,
г. Белая Церковь, Украина*

The aim of the study was to identify a valuable source material among available tomato sterile forms for tomato breeding for cold resistance.

Materials and methods. The article presents the results of a study of 14 tomato sterile forms for adaptive ability based on a complex trait of cold resistance, the elements of which were determined in field and laboratory conditions. The material was assessed according to a set of features: the increase in plant height, the rate of growth recovery, the duration of the sowing-seedling and seedling-fruit ripening periods. Calculations of the parameters of general adaptive ability, homeostatic and variance of specific adaptive ability of sterile form genotypes by earliness and cold resistance have been performed.

Results. The level of general and specific adaptive ability, as well as homeostatic of cold resistance, were determined. Valuable source material has been isolated according to the characteristics of cold resistance in the field and laboratory conditions.

The forms *Mar ps 59*, *Mar ps 53* and *Sat ps 52* were characterized by highest cold resistance in the laboratory conditions. These lines were characterized by seed germination in the range of 0,3–2,8 %, compared with other sterile forms with the germination 0,1–0,2 %. The increase in plant height under cool conditions in mentioned forms (*Mar ps 59*, *Mar ps 53* and *Sat ps 52*) was higher (45–54 %) compared with other sterile forms.

In the field, the duration of the interphase period «sowing – seedlings» was the shortest in the forms *Mar ps 59* and *Sat ps 52* compared with other forms studied. Thus, the duration of the period «sowing–germination» in the mentioned forms was 19 and 18 days respectively, against the standard *Flora 20* days. The increase in seedling height was 60–80 %, to the same extent with other forms and the standard 74 %.

Conclusions. The selected methods for assessing the traits of cold resistance and earliness and mathematical processing of the results allowed us to identify a valuable source material among sterile forms of tomatoes with a high level of adaptability of these traits.

Keywords: tomato, cold resistance, earliness, general and specific adaptive ability.

Most tomatoes are plants of the warm season and should be planted only after the danger of frost has passed. Tomato tolerance to low extreme temperatures is extremely important for the flowering phase and further fruit development. Flowering in the spring will occur if the daytime temperature is warm and the nighttime temperature does not drop below 13 °C. The temperature of growing tomato seedlings should be maintained at a constant level between 14–16 °C. Seedlings should not be transplanted until the last frosts [1].

Studies to increase the cold resistance of tomatoes using synthetic substances have shown that plant growth regulators reduce the negative effects of low temperatures by increasing the activity of cell membranes and reducing oxidative processes [2].

Complex crosses also contribute to the expansion of the genetic diversity of the original populations. At complex crosses selections on recessive signs can be already begun in the first hybrid generation. The use of complex crosses contributed to the origination of precocious, cold-resistant initial forms with valuable recessive traits, thus varieties *Zoren*, *Flora*, *Boyan* were created [3].

During normal growth and development, plants are exposed to various types of stress, such as drought, heat, ultraviolet light, air pollution and pathogen attack. Most plants receive physiological and biochemical damage due to exposure to temperatures above or below optimal for growth and development. As a result, these injuries are reflected in most metabolic processes, which reduces the ability to grow crops and, consequently, commercial yields [4].

Sterile forms of tomato are used to create heterosis hybrids of the first generation. At the current level of breeding due to the need to reduce the complexity of the hybridization process, the need for sterile forms of tomatoes is growing. The use of male sterility in tomato plants does not require emasculation of flowers

and allows to improve and reduce the cost of hybrid seed production.

In tomato plants, several types of sterility are known due to the *ms*, *sl*, *ps* genes: longostilia, functional male sterility (FMS), pollen genetic sterility, stamen sterility, stamenlessness of flowers [5]. The most promising and suitable for mass production of hybrid seeds is FMS caused by the *ps* gene [6].

In our studies we used the phenomenon of functional male sterility (FMS), which is due to morphological deviations from the normal structure of the flower (*ps* gene) [7–9]. To create F₁ hybrids we studied tomato samples with FMS which is characterized by the inability to crack anthers and is controlled by the *ps* gene.

Using the variety John Bear (donor of the *ps* gene) at the Kyiv research station, breeder V.A. Kravchenko created breeding sterile lines using sterility (*ps* gene): Tat *ps*, Kyiv *ps*, Mar *ps*, Sat *ps* and others. Plants of such lines formed 1–2 small fruits on a plant without seeds during the growing season. Compared to other types of sterility the reproduction of such sterile lines is simple.

Cold resistance of plants at the beginning of the growing season and during the functioning of the reproductive organs is important for breeding for heterosis. When creating cold-resistant and early ripening F₁ hybrids, the ability to grow rapidly at low temperatures in the cool early spring period is important. This study is devoted to the identification of valuable source material with good cold resistance traits among available sterile forms.

Material and Methods

The research was carried out for 2016-2018 years on the experimental field of Bila Tserkva National Agrarian University in the collection nursery. We studied 14 sterile forms of tomato obtained from the Kyiv research station and created by Professor V.A. Kravchenko.

Establishment of experiments, study and evaluation of sterile forms were carried out in accordance with the [10]. The study was conducted in comparison with the standard Flora variety. Cold resistance was determined by the methods [11, 12].

Calculations of the parameters of general adaptive ability, homeostatic and variance of specific adaptive ability of sterile form genotypes by earliness and cold resistance were performed according to the method of [13, 14]. The obtained results were processed according to the [15].

Results

The complex trait of cold resistance of sterile forms was evaluated taking into account simple traits that are easy to assess: the number of germinating seeds, the increase in plant height under cold stress and the rate of resumption when returning to favourable conditions in laboratory and field experiments. The forms Mar *ps* 59, Mar *ps* 53 and Sat *ps* 52 were characterized by highest cold resistance in the laboratory conditions (Table 1). These lines were characterized by seed germination in the range of 0,3–2,8 %, compared with other sterile forms with the germination 0,1–0,2 %. The increase in plant height under cool conditions in mentioned forms (Mar *ps* 59, Mar *ps* 53 and Sat *ps* 52) was higher (45–54 %) compared with other sterile forms. The plants quickly recovered growth in the early ontogenesis stages after exposure to low temperatures (10–13 %) at the level of the standard variety Flora (10 %).

Table 1.

Evaluation of the traits forming the cold resistance of tomato sterile forms in the laboratory (average for 3 years)

Variety, line	Trait											
	Seed germination				Increase of plant height				Growth resumption rate			
	%	Hom	GAAi	σSAAi	%	Hom	GAAi	σSAAi	%	Hom	GAAi	σSAAi
Flora – standard	6,0	0,06	0,0	1,7	50	0,54	0,1	4,0	10	0,11	0,0	16,1
Sat <i>ps</i> 51	0,5	0,01	-0,1	0,3	43	0,95	-0,1	2,7	9	0,15	2,3	7,0
Mar <i>ps</i> 53	2,0	0,03	0,2	0,7	54	0,66	0,6	4,0	10	0,11	-4,0	1,3
Mar <i>ps</i> 59	2,8	0,03	0,6	1,3	48	1,17	0,9	3,7	12	0,31	-3,4	0,7

Sat <i>ps</i> 52	0,3	0,01	-0,1	0,1	45	0,85	-0,1	2,7	13	0,46	0,2	0,5
LSD ₀₅	2,1				14,4				4,5			

Note: GAAi – general adaptive ability; Hom – homeostatic; σ SAAi – variance of specific adaptability; LSD₀₅ – least significant difference at 0,05 significance level

In the laboratory the sterile form Mar *ps* 59 was characterized by the highest homeostatic among sterile formson the trait «seed germination» 0,03, and on the trait «increase in plant height» 1,17, compared to the standard 0,06 and 0,54 respectively, and on the trait «growth resumption rate» 0,31 (0,11 in the standard).

The adaptive ability of hybrids is a valuable feature in selection for heterosis, so the creation of original sterile forms with high adaptive ability is important for breeding and seed production of tomato heterosis hybrids F₁. Under conditions of reduced variable temperatures in the laboratory on the basis of traits «seed germination» and «increase of plant height», we identified sterile forms of Mar *ps* 53 and Mar *ps* 59 with general adaptive ability (GAAi) in the range of 0,2–0,9and variance of specific adaptive ability (σ SAAi) in the range of 0,7–4,0 (Table 1).

The sterile form Mar *ps* 58 differed from other samples by the highest level of specific adaptive ability on the basis of the «growth resumption rate» of plants in the laboratory, which was 10,3.

The period from sowing to germination is important to determine early maturity and cold resistance. Tomato seeds of sterile forms showed different terms of germination under conditions of low temperatures. Under changing environmental conditions, resistance or instability to cold in the early stages of ontogenesis may not be maintained at the stage of functioning of the reproductive system. Therefore, the definition of cold resistance under the action of low temperatures on the growth and development of seeds and seedlings may not fully reveal the complex cold resistance [12, 16, 17].

To further test for cold tolerance we planted the seedlings to be exposed to periodic early spring cold snaps. When the temperature decreased to – 2 °C, sterile forms of Mar *ps* 59 and Sat *ps* 52 were characterized by high cold resistance (4,8 and 4,3 points respectivelyon a 5-point scalein accordance with [10] (Table 2).

Table 2.

Homeostatic and adaptive ability of cold resistance traits of tomato sterile forms in the field (average for 3 years)

Variety, line	Sowing - seedlings period				Trait							
					Increase of plant height				Cold resistance of seedlings			
	Days	Hom	GAAi	σ SAAi	%	Hom	GAAi	σ SAAi	Point	Hom	GAAi	σ SAAi
Flora – standard	20	0,67	5,4	0,7	74	1,84	5,0	0,5	2,5	0,09	0,8	0,2
Sat <i>ps</i> 51	21	0,67	0,6	5,3	80	2,84	-1,3	0,0	4,5	0,29	0,4	0,5
Mar <i>ps</i> 53	19	0,40	-0,1	6,0	55	2,14	0,7	1,0	4,0	0,49	-0,1	0,0
Mar <i>ps</i> 59	19	0,40	0,6	7,3	60	1,27	0,7	1,0	4,8	0,19	0,4	0,5
Sat <i>ps</i> 52	18	0,55	1,2	4,7	80	1,41	-1,3	0,0	4,3	0,17	0,2	0,2
LSD ₀₅	1,8				8,6				1,1			

Note: GAAi – general adaptive ability; Hom – homeostatic; σ SAAi – variance of specific adaptability; LSD₀₅ – least significant difference at 0,05 significance level

In the field, the duration of the interphase period «sowing – seedlings» was the shortest in the forms

Mar *ps* 59 and Sat *ps* 52 compared with other forms studied. Thus, the duration of the period «sowing – germination» in the mentioned forms was 19 and 18 days respectively, against the standard Flora, 20 days. The increase in seedling height was 60–80 %, to the same extent with other forms and the standard, 74 %.

In the field, sterile forms Mar *ps* 53, Sat *ps* 51 were isolated, which had a homeostatic index 0,40 and 0,67 respectively for the period «sowing – seedlings», and 2,14 and 2,84 for the «increase of plant height» (1,84 in the standard Flora). These lines had high homeostatic on the trait of «cold resistance of seedlings», 0,29 and 0,49 compared to the standard, 0,09, and other samples (Table 2). Different meanings in the laboratory and in the field, in our opinion, indicate the dependence of the trait expression on the variability of weather conditions over the years of cultivation.

In the field, the highest level of adaptability was observed in the sterile form Mar *ps* 59 for both general and specific adaptive ability. The value of GAA_i was 0,4–0,7 for all three traits, the value of σSAA_i was 0,5–7,3 for all three traits.

Conclusions

Analysing the traits of cold resistance, we found that sterile forms Mar *ps* 59 and Sat *ps* 52 have the highest performance in laboratory and field conditions and are the best in terms of cold resistance among the studied lines. This character is valuable for use in breeding for precocious in tomatoes. To create early ripening, cold-resistant heterosis F_1 hybrids, we recommend using them as source forms.

References

1. Grant A. (2020) Tomato Temperature Tolerance: Best Growing Temp For Tomatoes. Gardening Know How. Available at: <https://www.gardeningknowhow.com/edible/vegetables/tomato/growing-temp-for-tomatoes.htm>
2. Astakhova N.V., Suvorov T.A., Deryabina A.N., Trunov T.I. (2010) Influence of the drug Amerol-2000 on morphological parameters and cold resistance of tomato plants. *Agrokhimiya*, 2, 21-25.
3. Kravchenko V.A. (2014) Methodical approaches to the selection process in vegetable growing. *Novitni Ahrotekhnolohii*, 1 (2), 42-48.
4. Rivero R.M., Ruiz J.M., García P.C., López-Lefebvre L.R., Sánchez E., Romero L. (2001) Resistance to cold and heat stress: accumulation of phenolic compounds in tomato and watermelon plants. *Plant Science*, 160 (2), 315-321. doi: 10.1016/s0168-9452(00)00395-2.
5. Uzun I.V. (2016) Development of tomato hybrids based on forms of female parents with functional male sterility. *Vegetable crops of Russia*, 1, 24-28. In Russian. <https://doi.org/10.18619/2072-9146-2016-1-24-28>
6. Miroschnichenko T.M., Ivchenko T.V. (2013) Thereproduction of sterile forms of tomato for heterosis breeding using biotechnology techniques. *Ovovichnytstvo i bashtannytstvo*, 59, 212-218.
7. Kravchenko V.A., Kuzemsky A.V. and Ermakova I.V. 1998. Use of functional male sterility in the creation of heterotic tomato hybrids. *Vegetable and melon growing, Ukraine*, 43, 81-87.
8. Simonov A.A. (1967). Types of sterile tomatoes and their use for obtaining hybrid seeds. *Bulletin of Agricultural Sciences, Moscow*, 2: 27-33.
9. Atanassova B. (1999). Functional male sterility (*ps-2*) in tomato and its application in breeding and hybrid seed production. *Euphytica*, 107 (1), 13–21. <https://doi.org/10.1023/A:1003527714805>
10. Methodology of experimental work in vegetable and melon growing. (1992). Moscow: Agropromizdat.
11. Diagnostics of plant resistance to stress. Methodical guidance. (1988). Leningrad (pp. 75–227).
12. Polesskaya L.M., Zhakote A.G., German M.E. and Kharti V.G. (1991). Diagnostic traits of tomato resistance to low temperatures. *Bulletin of the Academy of Sciences of the Moldavian SSR. Biol. and Chem. Science*, 2, 20-23.
13. Kilchevskiy A. & Khotyleva L. (1985). Methods of assessments of the adaptive abilities and stability of genotypes differentiating the abilities of environment. *Messages 1. Justification of the method. Genetics, USSR*, 21(9), 1481-1490.
14. Kilchevskiy A. V. & Khotyleva L.V. (1985). Methods of assessments of the adaptive abilities and stabilities of genotypes differentiating abilities of the environment. 2. Numerical example and discussion. *Genetics, USSR*, 21(9), 1491-1498.
15. Guidelines for the mathematical processing of the results of counts and observations in breeding and genetic research. (1979). Moscow: Kolos.
16. Polonskiy V.I. & Kalinina L.M. (1990). Method for determining the cold resistance of tomatoes. *Bulletin of agricultural science, Moscow*, 8, 139 - 143.
17. Tsaranu L.A. & Tsaranu M.Kh. (2002). Assessment of the variability of the cold resistance trait in mutants of tomato in a homo- and heterozygous state. *Vegetable and melon growing, Ukraine*, 47, 19-25.