



UDC 636.2.034

DOI: 10.48077/scihor.25(8).2022.59-66

Relationship Between Globulins in the Late Dry Period with Biochemical Parameters, Fertility and Culling of Cows within 90 Days after Calving

Apollinariy Kraevskiy¹, Valentyn Yefimov², Vasyl Stefanyk^{3*}, Svitlana Vlasenko⁴, Taras Basarab³

¹Volodymyr Dahl East Ukrainian National University
49050, 12 Naukova Str., Dnipro, Ukraine

²Dnipro State Agrarian and Economic University
49600, 25 S. Yefremov Str., Dnipro, Ukraine

³Stepan Gzhytskyi National University of Veterinary Medicine and Biotechnologies Lviv
29010, 50 Pekarska Str., Lviv, Ukraine

⁴Bila Tserkva National Agrarian University
09117, 8/1 Soborna Sq., Bila Tserkva, Ukraine

Article's History:

Received: 11.08.2022

Revised: 22.09.2022

Accepted: 17.10.2022

Suggested Citation:

Kraevskiy, A., Yefimov, V., Stefanyk, V., Vlasenko, S., & Basarab, T. (2022). Relationship between globulins in the late dry period with biochemical parameters, fertility and culling of cows within 90 days after calving. *Scientific Horizons*, 25(8), 59-66.

Abstract. Diseases of the transition period in cows and their culling after calving attract considerable attention. Therefore, early diagnosis and prediction of pathological processes during the transition period in cows using biochemical markers stays relevant. The purpose of this study was to find the relationship of globulin levels in the late transition period with biochemical parameters and the level of fertilisation and culling of cows within 90 days after calving. The blood serum of cows of the main herd was investigated 10-15 days before calving, based on which the animals were conditionally divided into three groups (with signs of hypogammaglobulinemia (less than 30 g/l)), as well as with normal and elevated (more than 40 g/l) globulin levels. Apart from indicators of protein-nitrogen metabolism, enzymatic activity was also found in the blood serum and the state of carbohydrate-lipid, mineral, and vitamin metabolism was assessed using biochemical, chromatographic, and spectrophotometric research methods. It was found that cows with hypoglobulinemia showed substantially lower activity of asparagine transaminase, as well as differences in carbohydrate-lipid metabolism (high glucose levels and reduced concentration of high-density lipoproteins). Furthermore, these cows were found to have a decrease in the content of magnesium, as well as copper. It was discovered that within 90 days after calving, the highest level of culling was observed among cows with reduced globulin levels, and the lowest – in animals with normoglobulinemia. Therefore, a direct relationship between the level of globulins in the blood serum 10-15 days before calving with the reproductive capacity and the level of culling of cows was proved. This allows promptly diagnosing and predicting the development of post-partum pathology and reproductive disorders. The results obtained can be used by practical specialists and scientists to plan and investigate the effectiveness of preventive measures to improve the safety of cows and their reproductive capacity in the post-calving period

Keywords: transition period, post-partum pathology, globulins, biochemical parameters, mineral metabolism, reproductive ability



Copyright © The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (<https://creativecommons.org/licenses/by/4.0/>)

*Corresponding author

INTRODUCTION

In dairy cows, transition diseases lead to considerable economic losses in the dairy industry. These losses include the cost of treatment and premature culling of cows. Early diagnosis or prediction of transition pathologies allows developing measures for their prevention, which reduces the incidence rate and reduces economic losses in the dairy industry (Venjakob *et al.*, 2021; Hailemariam *et al.*, 2014).

In the conditions of industrial cattle breeding and high milk productivity, the duration of the reproductive life of cows does not exceed 2.5-3 lactations. One of the reasons for this is a violation of reproductive function, caused by the wide spread of diseases of the reproductive system in the prenatal, birth, and post-partum periods (Yazlik *et al.*, 2020; Sewalem *et al.*, 2008). The rate of culling cows from the herd on dairy farms during the year is about 21.0-38.2% (Pinedo *et al.*, 2010; Stojic *et al.*, 2013). According to Diniso & Jaja (2021), the most common causes of this are reproductive issues, namely infertility (7.9%), reproductive disorders (89.9%) and dystocia (1%). At the same time, successful fertilisation of cows in the post-partum period is crucial for further reproductive efficiency and profitability in modern dairy cattle breeding (Sammad *et al.*, 2022).

Many post-partum diseases of cows are based on a violation of metabolic processes during the transition period, which determine the further condition of animals in the post-natal period, their reproductive ability and the level of culling of animals. The development of energy and protein deficiencies leads to a decrease in the activity of polymorphonuclear neutrophils (Yazlik *et al.*, 2020), as well as the release of inflammatory cytokines (tumour necrosis factor α and interleukin-6), which lead to systemic inflammation and insulin resistance (Wankhade *et al.*, 2017). Systemic inflammation in the early post-partum period is negatively associated with the ovulatory status of the first dominant follicle after calving (Cheong, 2017).

During the transition period, there are substantial changes in the functioning of organs and the hormonal status of the animal body (Kucuk Baykan & Özcan, 2019). Cows' appetite worsens, and the amount of feed eaten does not compensate for the rapidly increasing cost of milk production. Animals compensate for this lack by mobilising the reserves of fats, proteins, micro- and macronutrients of their body. It is no coincidence that it is during this period that dystrophy and cirrhosis of the liver, emaciation (cachexia), rumen acidosis, ketosis, inflammation of the reproductive organs and, as a result, culling and loss of animals take place (Grummer, 1995). The situation is aggravated by the development of hypocalcaemia and the intensification of protein metabolism in the post-partum period, which has adverse consequences for the health and reproductive ability of dairy cows (Sammad *et al.*, 2022).

Violations of metabolic processes occurring in the body are primarily accompanied by changes in several

biochemical parameters. In this regard, the search for individual proteins, metabolites, enzymes, and hormones that would have diagnostic and prognostic significance continues. Andjelić *et al.* (2022) show that higher serum calcium concentrations after calving are associated with higher levels of progesterone, glucose, and triglycerides. In turn, subclinical hypocalcaemia leads to the development of metritis, prolongation of the service period and a decrease in the size of the follicle (Paiano *et al.*, 2020).

Blood serum proteins are also related to the physiological condition of cows. As shown by Stravsky *et al.* (2020), the albumin-globulin ratio can be used as a prognostic factor to identify cows prone to uterine subinvolution. Globulin concentrations decrease 5 weeks before calving and increase after calving, and their change during calving is substantially related to the frequency of fertilisation (Rowlands *et al.*, 1980).

The purpose of this study was to find the level of culling and fertilisation capacity of cows within 90 days after calving, depending on the level of globulins and other biochemical parameters in the blood serum in the late transition period.

MATERIALS AND METHODS

Selection of animals and organisation of the experiment.

The experimental part of the studies was performed on 37 dairy cows of 4-7 years of age with an average annual productivity of 7,200 kg of milk on one of the commercial farms of the Astarta-Kyiv agro-industrial holding. Biochemical studies were performed 10-15 days before calving. According to the results of the study, cows were conditionally divided into three groups. The first group (5 animals) included cows with a low globulin level of 25.6 ± 0.8 g/l (in all animals – up to 30 g/l). The second (control) group (10 cows) was formed from animals with an average globulin content of 35.3 ± 0.9 g/l (within 30-40 g/l), which corresponded to the level of conditionally normal indicators, and the third group included 22 cows with a globulin content of 49.9 ± 1.5 g/l (within 40-65 g/l). Thus, according to the level of globulins, the second group was assigned to conditionally healthy animals (the first group included cows with hypogammaglobunemia, and the third group included cows with hypergammaglobunemia).

The general condition of the cows was monitored for 90 days after calving. Therewith, the number of culled cows was recorded due to obstetric pathology, mastitis, rennet displacement, and diseases of the distal extremities. Furthermore, fertilisation and infertility rates were identified in animals of each group.

Sampling. Blood for biochemical studies was taken 10-15 days before calving from the tail vein before feeding the animals. Selected blood samples were used to obtain a serum, which was frozen at -20°C and transported and stored in this state until the time of research. Biochemical studies were carried out under

the conditions of the laboratory of clinical biochemistry of the Research Centre for Biosafety and Environmental Control of Agricultural Resources of the Dnipro State Agrarian and Economic University.

Biochemical studies. The blood serum samples were examined for the content of total protein (with biuret reagent), albumin (with bromocresol green), urea (enzymatically according to the Berthelot reaction), creatinine (by the rate of formation of the creatinine-picrate complex in the Jaffe reaction), glucose (glucose oxidase method), total calcium (with arsenazo III), inorganic phosphorus (with molybdenum reagent), potassium (by turbidimetric method with sodium tetraphenylborate), iron (by reaction with ferrozine). An automatic biochemical analyser Miura-200 (Italy) was used for analysis with ready-made sets of reagents produced by Spinreakt (Spain), Dialab (Austria), Cormay (Poland), and HTI (USA). High-density lipoproteins were determined according to Warnick & Albers (1978).

Sodium levels were found by colorimetric reaction with magnesium-uranyl acetate (Human, Germany), and magnesium – by colour reaction with calmagite (HTI, USA) on a semi-automatic biochemical analyser Humalazer 3000 (Human, Germany).

The activity of transamination enzymes (ALT and AST) was found kinetically using a set of Spinreakt reagents (Spain), alkaline phosphatase – by the rate of 4-nitrophenol formation (Cormay, Poland). The carotene concentration was found spectrophotometrically (Ulab-2, China) after precipitation of serum proteins with ethyl alcohol and extraction of carotene with hexane.

The concentration of vitamins A and E was found using high-performance liquid chromatography on an Agilent Technologies 1260 Infinity liquid chromatograph, column C18 with spectrophotometric detection (Yefimov & Sofonova, 2014). Reagents manufactured by Honeywell (USA) and Carlo Erba (Italy) were used to manufacture the components of the solvent system.

The concentration of copper, zinc, and manganese in blood serum samples was found by Atomic Absorption Spectrophotometry with flame atomisation on a Selmi FCM 115 spectrophotometer (Ukraine).

The globulin content and A/G ratio were found by calculation.

Statistical analysis. In each group of cows, statistical processing of all the biochemical parameters under study was performed. The resulting digital material was processed using variational statistics methods using SPSS Data editor 17.0 version according to the Tukey Test with Bonferroni correction.

RESULTS AND DISCUSSION

When analysing the results of biochemical studies, it was found that the total protein in the peripheral blood serum of cows at the end of the transition period depended mainly on the concentration of globulins, since the level of albumins practically did not change.

With a low level of globulins (the first group), the level of total protein in the blood serum was also 10.5% lower compared to animals with their physiological values (the second group), and 1.4 times lower compared to animals of the third group (Table 1).

Table 1. Biochemical parameters of blood serum of cows with different levels of globulins (M±m)

Indicators	Group 1	Group 2	Group 3
Total protein, g/l	57.4±2.542	67.9±2.2**	82.1±2.1***
Albumins, g/l	31.8±2.3	32.6±2.1	32.3±1.0
Globulins, g/l	25.6±0.8	35.3±0.9***	49.9±1.5***
A/G ratio, units	1.24±0.09	0.93±0.06*	0.65±0.02***
Creatinine, µmol/l	110.8±9.5	113.0±5.0	119.6±4.3
AST, IU/l	55.6±6.8	104.2±12.2**	91.2±7.8**
ALT, IU/l	19.4±1.9	17.6±1.9	19.7±1.5
Alkaline phosphatase, IU/l	96.5±8.0	109.0±12.2	101.3±6.2
Glucose, mmol/l	2.68±0.11	2.30±0.06**	2.30±0.05**
High-density lipoproteins, mg/100 ml	773.2±95.1	1,088.1±105.9*	1,073.0±76.2*

Note: * – $p \leq 0.05$; ** – $p \leq 0.01$; *** – $p \leq 0.001$ – relative to animals of the first group

Source: compiled by the authors

Accordingly, the A/G ratio was highest in cows of the first group and amounted to 1.24±0.09, in the second group it was lower by 33.3% ($p \leq 0.02$), and in the third – by 1.9 times ($p \leq 0.001$).

Apart from the above data on the indicators of total protein, globulins and albumins, certain differences in

the activity of transamination enzymes were noted. The activity of AST, which is an indicator enzyme for the liver in cattle, in animals of the first group was almost 2 times lower compared to the second and third groups ($p \leq 0.01$) and by 64.0% ($p \leq 0.01$), respectively. Indicators of carbohydrate-lipid metabolism in cows of the first group were

described by a high level of glucose and, at the same time, a low content of lipoproteins. The concentration of glucose in the blood of cows of the first group was 1.17 times higher ($p \leq 0.01$), and lipoproteins – 1.4 times ($P \leq 0.05$) lower than in animals of the second and third

groups. Considering the importance of indicators and the nature of mineral and vitamin metabolism in the pathogenesis of various pathologies of the transition period, the level of basic macro- and microelements, as well as vitamins A, E and carotene was investigated (Table 2).

Table 2. Indicators of mineral and vitamin metabolism in cows in the transition period with different levels of globulins ($M \pm m$)

Indicators	Group 1	Group 2	Group 3
Total calcium, mmol/l	2.40±0.14	2.13±0.16	2.25±0.10
Inorganic phosphorus, mmol/l	1.54±0.15	1.86±0.16	2.04±0.10*
Magnesium, mmol/l	0.92±0.02	2.02±0.66	1.53±0.31°
Potassium, mmol/l	5.02±0.78	4.53±0.62	4.98±0.40
Sodium, mmol/l	137.6±8.3	136.9±2.9	137.2±3.2
Iron, µmol/l	28.2±2.3	25.1±1.7	23.3±1.6
Copper, µg/100 ml	55.4±4.2	74.3±3.5**	82.5±3.0***
Zinc, µg/100 ml	76.0±4.5	76.5±7.3	83.8±4.7
Manganese, µg/100 ml	2.4±1.2	1.6±0.2	1.8±0.2
Vitamin A, µg/100 ml	18.3±0.6	19.1±2.6	19.3±1.0
Vitamin E, µg/ml	2.2±0.4	2.9±0.5	3.1±0.3°
Carotene, µg/100 ml	306.2±11.6	261.2±15.4*	287.3±10.9

Note: ° $p \leq 0.1$; * – $p \leq 0.05$; ** – $p \leq 0.01$; *** – $p \leq 0.001$ – relative to animals of the first group

Source: compiled by the authors

Considering the indicators of mineral metabolism, one should pay attention to the level of calcium, phosphorus, and its ratio in the blood serum of cows with various levels of globulins. The total calcium content in animals of separate groups did not significantly differ. At the same time, animals of the second and third groups showed an increase in the level of inorganic phosphorus, which was significant in cows with a globulin content of more than 40 g/L. A violation of mineral metabolism is indicated by a tendency to reduce the level

of magnesium in the blood of cows of the first group. Copper levels were 1.3 ($p \leq 0.01$) and 1.5 ($p \leq 0.001$) times lower than in animals of the second and third groups, respectively.

The content of other macro- and microelements in the blood serum of cows of all groups did not differ. The frequency of culling and the results of the diagnostic stage of obstetric and gynaecological medical examination in all groups of cows during 2-3 months of lactation are presented in Table 3.

Table 3. Results of obstetric and gynaecological medical examination of cows within 90 days after calving

Indicators	First group		Second group		Third group	
	n	%	n	%	n	%
Culled	4	80.0	1	10.0	5	22.7
Pregnant	0	0	7	70.0	5	22.7
Barren	1	20.0	2	20.0	12	54.6

Source: compiled by the authors

Within 2-3 months after calving, 4 cows (80%) were culled from the herd in the first group, which is significantly more than in the second group – 1 cow (10%) and third group – 5 cows (22.7%). The reasons for culling were obstetric and gynaecological pathology (development of chronic mastitis and/or displacement of the rennet), as well as diseases of the distal extremities. At the same time, the metabolic status of cows in the second and third groups was affected during the

post-partum period and reproductive function during the experiment. The fertilisation rate of cows of different groups within 3 months after calving also differed. In cows of the second group, it was 70.0%, which is 47.3% higher than in the third group, and the number of infertile cows was 34.6% higher in the third group.

Studies have shown that before calving, the majority of cows (73%) had globulin levels higher or lower than animals with conditionally normal values (Brcsic

et al., 2015; Alberghina et al., 2011; Kahn et al., 2010). It is known that among the protein globulin fractions, gamma globulins predominate, which provide the function of immune protection of animals (Elshahawy & Abdullaziz, 2017; Bertoni et al., 2008). Therefore, the low concentration of these proteins in cows of the first group has a low level of their nonspecific resistance. At the same time, elevated globulin levels in cows of the third group may be due to increased protein levels during the acute phase of inflammation. As noted by Tothova et al. (2016), they belong to the α -fraction, and their level increases when there is a focus of inflammation or a systemic inflammatory response in the body. That is, the noted changes may indicate the activation of the immune system and the activation of inflammatory processes before calving, including in the liver. This is indicated by the research of Bertoni et al. (2008), who established the dependence of the level of α -globulins in the first weeks after childbirth and the frequency of violations of reproductive ability in cows.

In animals with reference globulin levels, the concentration of total protein and albumins also did not exceed normal values. Notably, the concentration of albumins did not differ between animals of different groups. According to Spaans et al. (2022), the globulin concentration decreases 7-14 days before calving, but the albumin content remains relatively stable.

Changes in the content of globulins in the blood serum are probably closely related to general disorders of protein metabolism. Cows with low levels of globulin protein fractions also had low levels of total protein. On the one hand, this may be caused by the high intensity of the use of the total pool of amino acids in the mother's blood for protein synthesis in the foetal body. At the same time, this may indicate the use of immunoglobulins G for colostrum formation during the transient period (Conneely et al., 2013).

Ingvartsen (2006) indicates that AST levels in healthy cows range from 78 to 132 IU/L. Its increase can be observed at the end of the transition period, since uterine tissues have a fairly high activity of this enzyme (Sattler & Fürll, 2004). Presumably, a decrease in AST activity in cows against the background of a decrease in globulin levels may indicate a decrease in the number of functionally active hepatocytes due to the previous cytolytic syndrome and replacement of parenchymal cells with connective tissue. Thus, the hepatocyte resource of the liver is depleted with damage to hepatocytes because of autointoxication with decay products and xenobiotics (Djoković et al., 2013).

Changes in carbohydrate-lipid metabolism indicate that cows of the second and third groups had a lower glucose level against the background of an increase in the concentration of high-density lipoproteins in their blood serum. Such biochemical changes are inherent in the development of a negative energy balance, which begins to develop in highly productive cows at the end

of the transient period (Ha et al., 2022). At the same time, blood glucose levels in cows with low globulin levels were higher than in animals of the second and third groups. The content of lipoproteins, on the contrary, was lower. Such dynamics of indicators can be explained by a lower intensity of colostrum formation and, accordingly, lower requirements for metabolic energy for its production.

Analysing the reduced copper content, Kaya et al. (2016) indicates that 95% of the copper in serum or plasma is part of the protein ceruloplasmin, which is synthesised in the liver. Therefore, it can be assumed that it is functional liver failure that causes a decrease in copper concentration. Nobili et al. (2013) noted a decrease in ceruloplasmin activity in children with non-alcoholic fatty hepatodystrophy. Such changes lead both to a decrease in neutrophil activity in cows and to a decrease in the level of antioxidant protection, which depends on the level of ceruloplasmin (Hernández et al., 2022).

The obtained research results coincide with the data of Rowlands et al. (1980) who indicate that pre-calving globulin concentrations are related to subsequent reproductive performance of cows. Thus, at the end of the transition period, changes in metabolic processes and the development of immunodeficiency in cows with hypoglobulinemia are noted. Among these animals, 80% were subsequently culled within three months of calving. Apparently, cows with weakened immunity, especially after calving, are more prone to many diseases of the transient period, which is also indicated by Zhelavskiy et al. (2020). As a result of culling up to 28.0% in the first 100 days after calving (Kucuk Baykan & Özcan, 2019). Therewith, the reasons for culling of animals from the herd are different – from ketosis to rennet displacement, mastitis, limb pathology, which coincides with the results obtained.

CONCLUSIONS

Given that the limits of globulin levels did not overlap between the groups of animals under study, it became possible to identify pathognomic indicators of metabolic disorders and use them to diagnose and predict the complicated course of the transient period, which leads to premature culling of animals. These pathological processes are accompanied by an increased or reduced content of globulins, due to the pathological condition of the liver and/or other organs that can be detected even before calving.

In the blood serum of cows with reduced globulin levels, AST activity decreases, glucose levels are increased, as well as lower concentrations of copper and lipoproteins, which indicates damage to the liver and other organs. As a result, their culling from the herd in the first 90 days after calving reaches 80%. At the same time, an increased level of blood globulins in the pre-pregnancy period is associated with an increased level of infertility, which reaches 54.6%. This

may indicate the development of a general inflammatory syndrome due to pathological processes in the organs of the reproductive system. In the blood serum of cows with reduced globulin levels, glucose was 1.17 times higher, while lipoproteins were 1.4 times lower than in the second and third groups. Cows with hypoglobulinemia develop mineral metabolism disorders, specifically a decrease in the level of magnesium and copper in the blood. Therefore, the level of globulins in the blood serum of cows in the late transition period

can be used as a prognostic indicator of the complicated course of the post-calving period. Globulin levels below 30 g/l 10-14 days before calving should be considered the greatest risk for the culling of cows. Exceeding the indicator over 40 g/l in the future significantly increases the frequency of reproductive function disorders.

Further studies may be aimed at evaluating the effectiveness of therapeutic and preventive measures in animals with detected violations of globulin levels in the late transition period.

REFERENCES

- [1] Alberghina, D., Giannetto, C., Vazzana, I., Ferrantelli, V., & Piccione, G. (2011). Reference intervals for total protein concentration, serum protein fractions, and albumin/globulin ratios in clinically healthy dairy cows. *Journal of Veterinary Diagnostic Investigation*, 23(1), 111-114. doi: 10.1177/104063871102300119.
- [2] Andjelić, B., Djoković, R., Cincović, M., Bogosavljević-Bošković, S., Petrović, M., Mladenović, J., & Čukić, A. (2022). Relationships between milk and blood biochemical parameters and metabolic status in dairy cows during lactation. *Metabolites*, 12(8), article number 733.
- [3] Bertoni, G., Trevisi, E., Han X., Bionaz, M., & Bionaz, M. (2008). Effects of inflammatory conditions on liver activity in puerperium period and consequences for performance in dairy cows. *Journal of Dairy Science*, 91(9), 3300-3310. doi: 10.3168/jds.2008-0995.
- [4] Brscic, M., Cozzi, G., Lora, I., Stefani, A.L., Contiero, B., Ravarotto, L., & Gottardo, F. (2015). Reference limits for blood analytes in Holstein late-pregnant heifers and dry cows: Effects of parity, days relative to calving, and season. *Journal of Dairy Science*, 98(11), 7886-7892. doi: 10.3168/jds.2015-9345.
- [5] Cheong, S.H., Sá Filho, O.G., Absalon-Medina, V.A., Schneider, A., Butler, W.R., & Gilbert, R.O. (2017). Uterine and systemic inflammation influences ovarian follicular function in postpartum dairy cows. *PLoS One*, 12(5), article number e0177356.
- [6] Conneely, M., Berry, D.P., Sayers, R., Murphy, J.P., Lorenz, I., Doherty, M.L., & Kennedy, E. (2013). Factors associated with the concentration of immunoglobulin G in the colostrum of dairy cows. *Animal*, 7(11), 1824-1832. doi: 10.1017/S1751731113001444.
- [7] Diniso, Y.S., & Jaja, I.F. (2021). A retrospective survey of the factors responsible for culling and mortality in dairy farms in the Eastern Cape Province, South Africa. *Scientific African*, 12, article number e00838.
- [8] Djoković, R., Šamanc, H., Jovanović, M., Fratrić, N., Dosković, V., & Stanimirović, Z. (2013). Relationship among blood indicators of hepatic function and lipid content in the liver during transitional period in high-yielding dairy cows. *Acta Scientiae Veterinariae*, 41(1), 1-6.
- [9] Elshahawy, I.I., & Abdullaziz, I.A. (2017). Hemato-biochemical profiling in relation to metabolic disorders in transition dairy cows. *Alexandria Journal for Veterinary Sciences*, 55(2), 25-33.
- [10] Grummer, R.R. (1995). Impact of changes in organic nutrient metabolism on feeding the transition dairy cow. *Journal of Animal Science*, 73, 2820-2833. doi: 10.2527/1995.7392820x.
- [11] Ha, S., Kang, S., Han, M., Lee, J., Chung, H., Oh, S.-I., Kim, S., & Park, J. (2022). Predicting ketosis during the transition period in Holstein Friesian cows using hematological and serum biochemical parameters on the calving date. *Scientific Reports*, 12(1), 1-9.
- [12] Hailemariam, D., Mandal, R., Saleem, F., Dunn, S.M., Wishart, D.S., & Ametaj, B.N. (2010). Identification of predictive biomarkers of disease state in transition dairy cows. *Journal of Dairy Science*, 93, 2680-2693. doi: 10.3168/jds.2010-6803.
- [13] Hernández, J., Muiño, R., Bedito, J.L., Abuelo, A., & Castillo, C. (2022). Redox status and oxidative stress in bovine. *Large Animal Review*, 28(3), 145-151.
- [14] Ingvarsen, K.L. (2006). Feeding-and management-related diseases in the transition cow: Physiological adaptations around calving and strategies to reduce feeding-related diseases. *Animal Feed Science and Technology*, 126(3-4), 175-213. doi: 10.1016/j.anifeedsci.2005.08.003.
- [15] Kahn, C.M., & Line, S. (Eds.) (2010). *The Merck veterinary manual*. New York: Merck & Co., Inc.
- [16] Kaya, S., Merhan, O., Kacar, C., Colak, A., & Bozukluhan, K. (2016). Determination of ceruloplasmin, some other acute phase proteins, and biochemical parameters in cows with endometritis. *Veterinary World*, 9(10), article number 1056. doi: 10.14202/vetworld.2016.1056-1062.

- [17] Kucuk Baykan, Z., & Özcan, M. (2019). Causes of culling and disease incidences at first production year of imported brown swiss and simmental cows from Austria. *Kocatepe Veterinary Journal*, 12(2), 178-184. doi: 10.30607/kvj.478070.
- [18] Nobili, V., Siotto, M., Bedogni, G., Ravà, L., Pietrobattista, A., Panera, N., Alisi, A., & Squitti, R. (2013). Levels of serum ceruloplasmin associate with pediatric nonalcoholic fatty liver disease. *Journal of Pediatric Gastroenterology and Nutrition*, 56(4), 370-375.
- [19] Paiano, R.B., Birgel, D.B., Bonilla, J., & Birgel Junior, E.H. (2020). Alterations in biochemical profiles and reproduction performance in postpartum dairy cows with metritis. *Reproduction in Domestic Animals*, 55(11), 1599-1606. doi: 10.1016/j.rvsc.2021.01.015.
- [20] Pinedo, P.J., De Vries, A., & Webb, D.W. (2010). Dynamics of culling risk with disposal codes reported by dairy herd improvement dairy herds. *Journal of Dairy Science*, 93, 2250-2261. doi: 10.3168/jds.2009-2572.
- [21] Rowlands, G.J., Manston, R., Stark, A.J., Russell, A.M., Collis, K.A., & Collis, S.C. (1980). Changes in albumin, globulin, glucose and cholesterol concentrations in the blood of dairy cows in late pregnancy and early lactation and relationships with subsequent fertility. *The Journal of Agricultural Science*, 94(3), 517-527. doi: 10.1017/S0021859600028501.
- [22] Sammad, A., Khan, M.Z., Abbas, Z., Hu, L., Ullah, Q., Wang, Y., Zhu, H., & Wang, Y. (2022). Major nutritional metabolic alterations influencing the reproductive system of postpartum dairy cows. *Metabolites*, 12(1), article number 60. doi: 10.3390/metabo12010060.
- [23] Sattler, T., & Fürll, M. (2004). Creatine kinase and aspartate aminotransferase in cows as indicators for endometritis. *Journal of Veterinary Medicine Series A*, 51(3), 132-137. doi: 10.1111/j.1439-0442.2004.00612.x.
- [24] Sewalem, A., Miglior, F., Kistemaker, G.J., Sullivan, P., & Van Doormaal, B.J. (2008). Relationship between reproduction traits and functional longevity in Canadian dairy cattle. *Journal of Dairy Science*, 91(4), 1660-1668. doi: 10.3168/jds.2007-0178.
- [25] Spaans, O.K., Kuhn-Sherlock, B., Hickey, A., Crookenden, M.A., Heiser, A., Burke, C.R., Phyn, C.V.C., & Roche, J.R. (2022). Temporal profiles describing markers of inflammation and metabolism during the transition period of pasture-based, seasonal-calving dairy cows. *Journal of Dairy Science*, 105(3), 2669-2698. doi: 10.3168/jds.2021-20883.
- [26] Stojic, P., Beskorovajni, R., Pantelic, V., Novakovic, Z., Kovacevic, S., & Stanojevic, D. (2013). Causes for culling first calving cows on farms with different levels of production. *Biotechnology in Animal Husbandry*, 29(2), 259-267. doi: 10.2298/BAH1302259S.
- [27] Stravsky, Y., Boltyk, N., Sachuk, R., Serheyev, V., & Rushchynska, T. (2020). The content of total protein and protein fractions in cows during pregnancy and their diagnostic value. *Scientific Messenger of LNU of Veterinary Medicine and Biotechnologies*, 22(99), 198-202. doi: 10.32718/nvlvet9930.
- [28] Tothova, C., Nagy, O., & Kovac, G. (2016). Serum proteins and their diagnostic utility in veterinary medicine: A review. *Veterinárni Medicína*, 61(9), 475-496. doi: 10.17221/19/2016-VETMED.
- [29] Venjakob, P.L., Staufenbiel, R., Heuwieser, W., & Borchardt, S. (2021). Association between serum calcium dynamics around parturition and common postpartum diseases in dairy cows. *Journal of Dairy Science*, 104(2), 2243-2253. doi: 10.3168/jds.2019-17821.
- [30] Wankhade, P.R., Manimaran, A., Kumaresan, A., Jeyakumar, S., Ramesha, K.P., Sejian, V., Rajendran, D., & Varghese, M.R. (2017). Metabolic and immunological changes in transition dairy cows. *Veterinary World*, 10(11), 1367-1377. doi: 10.14202/vetworld.2017.1367-1377.
- [31] Warnick, G.R., & Albers, J.J. (1978). A comprehensive evaluation of the heparin-manganese precipitation procedure for estimating high density lipoprotein cholesterol. *Journal of Lipid Research*, 19, 65-76.
- [32] Yazlik, M.O., Çolakoğlu, H.E., Kaya, U., Küplülü, Ş., & Vural, M.R. (2020). Does increased immune response at early postpartum period have a relationship with metabolic markers and subsequent fertility? *Polish Journal of Veterinary Sciences*, 23(1), 27-35. doi: 10.24425/pjvs.2020.132745.
- [33] Yefimov, V., & Sofonova, D. (2014). Age dynamics of concentrations vitamins a and e in the serum of piglets during a rearing period. *Science and Technology Bulletin of SRC for Biosafety and Environmental Control of AIC*, 2(3), 46-50.
- [34] Zhelavskiy, M.M., Kernychnyi, S.P., Mizyk, V.P., Dmytriv, O.Y., & Betlinska, T.V. (2020). The importance of metabolic processes and immune responses in the development of pathology of cows during pregnancy and postpartum periods. *Ukrainian Journal of Veterinary and Agricultural Sciences*, 3(2), 36-41. doi: 10.32718/ujvas3-2.06.

Взаємозв'язок рівня глобулінів у пізній сухостійний період з рівнем запліднення та вибуття корів протягом 90 діб після отелення

Аполінарій Йосипович Краєвський¹, Валентин Геннадійович Єфімов²,
Василь Юрійович Стефанік³, Світлана Анатолівна Власенко⁴, Тарас Петрович Басараб³

¹Східноукраїнський національний університет імені Володимира Даля
49050, вул. Наукова, 13, м. Дніпро, Україна

²Дніпровський державний аграрно-економічний університет
49600, вул. С. Єфремова, 25, м. Дніпро, Україна

³Львівський національний університет ветеринарної медицини та біотехнологій імені Степана Гжицького
79010, вул. пекарська, 50, м. Львів, Україна

⁴Білоцерківський національний аграрний університет
09117, Соборна площа, 8/1, м. Біла Церква, Україна

Анотація. Захворювання транзитного періоду у корів та їх вибуття після отелення привертають значну увагу. Тому рання діагностика та прогнозування патологічних процесів у перехідний період у корів із використанням біохімічних маркерів залишається актуальною. Метою роботи було визначити взаємозв'язок рівня глобулінів у пізній сухостійний період з біохімічними показниками та рівнем запліднення та вибуття корів протягом 90 діб після отелення. Було проведено дослідження сироватки крові корів основного стада за 10-15 діб до родів, на підставі чого умовно розділено тварин на три групи (з ознаками гіпогамаглобунемії (менше 30 г/л), а також з нормальним та підвищеним (понад 40 г/л) рівнем глобулінів. Крім показників білково-азотистого метаболізму, в сироватці крові також визначали ензиматичну активність та оцінювали стан вуглеводно-ліпідного, мінерального і вітамінного обмінів з використанням біохімічних, хроматографічних та спектрофотометричних методів досліджень. Було встановлено, що у корів з проявом гіпоглобулінемії спостерігалася суттєво нижча активність аспарагінової трансамінази, а також відмінності показників вуглеводно-ліпідного обміну (вищий рівень глюкози та знижена концентрація ліпопротеїдів високої щільності). Крім того, у цих корів було виявлено зниження вмісту магнію, а також міді. Досліджено, що впродовж 90 діб після отелення спостерігався найвищий рівень вибуття серед корів зі зниженим рівнем глобулінів, а найнижчий – у тварин з нормоглобулінемією. Отже, доведено безпосередній зв'язок між рівнем глобулінів у сироватці крові за 10–15 діб до отелення з репродуктивною здатністю та рівнем вибуття корів. Це дає змогу проводити ранню діагностику та прогнозування розвитку післяродової патології та порушення репродуктивної функції. Одержані результати можуть бути використані практичними фахівцями та науковцями для планування та вивчення ефективності профілактичних заходів з підвищення збереженості корів та їх відтворювальної здатності у післяотельний період.

Ключові слова: сухостійний період, післяродова патологія, глобуліни, біохімічні показники, мінеральний обмін, репродуктивна здатність